

# EXPERIMENT NO. 16

## Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations.**

You should indicate clearly at what stage in a test a change occurs.

Marks are **not** given for chemical equations.

**No additional tests for ions present should be attempted.**

**If any solution is warmed, a boiling tube MUST be used.**

Rinse and reuse test-tubes and boiling tubes where possible.

**Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.**

(a) **FA 6** is a solution of two different salts. It contains two different cations, one of which is listed in the Qualitative Analysis Notes. It contains two anions, both of which are listed in the Qualitative Analysis Notes.

- (i) Choose reagents that will allow you to identify one of the cations. Carry out suitable tests using these reagents and record your results in the space below.

I	
II	
III	
IV	
V	

One of the cations in **FA 6** is .....

(ii) Carry out the following tests to identify the two anions present in **FA 6**.

<i>test</i>	<i>observations</i>
To a 1 cm depth of <b>FA 6</b> in a test-tube add a 1 cm depth of aqueous silver nitrate, then	
add aqueous ammonia.	
To a 1 cm depth of <b>FA 6</b> in a test-tube add a 1 cm depth of aqueous barium chloride (or aqueous barium nitrate), then	
add dilute nitric acid.	

The anions in **FA 6** are ..... and .....

[9]

VI	
VII	
VIII	
IX	

(b) **FA 7** is an acidified solution of iron(II) sulfate,  $\text{FeSO}_4(\text{aq})$ .

Carry out the following tests and record your observations.

<i>test</i>	<i>observations</i>
(i) To a 1 cm depth of <b>FA 7</b> in a test-tube add aqueous sodium hydroxide and leave for a few minutes.	
(ii) To a 1 cm depth of <b>FA 7</b> in a boiling tube add a 1 cm depth of dilute sulfuric acid followed by a 1 cm depth of '20 vol' hydrogen peroxide. Stir the mixture, then	
(iii) pour a 1 cm depth of the mixture into a clean boiling tube and add a 3 cm depth of aqueous sodium hydroxide.	

I	
II	
III	
IV	
V	
VI	

(iv) What type of reaction takes place in (ii)?

.....  
.....

(v) Explain your observations in (iii).

.....  
.....

[6]

[Total: 15]